
Spine Injuries and Their Surgical Treatment

B. M. Isakov, K. B. Isakov
Andijan State Medical Institute

Abstract: injuries of the spine are a severe type of injury, accompanied in most cases by the disability of patients. Objective of the study: to improve the results of surgical treatment of injuries and diseases of the spine using tpf systems. The results of 114 patients with injuries of the thoracic and lumbar spine were analyzed. Mechanism of injury: fall from a height - 88 patients, road injury - 33 patients. Of the total number of examined patients, 53 patients underwent conservative treatment and 61 patients underwent surgical treatment. 1 fixation of the damaged segment of the spine with various types of metal structures (tpf) - 22 patients. 2 decompressive laminectomy with revision of the spinal canal and spinal cord at the level of injury (without fixation of the spinal column) – 20 patients. 3 decompressive laminectomy at the level of damage + revision of the epi- and subdural spaces of the spinal cord and, as the final stage of the operation, fixation with metal structures - 19 patients. Results: during the surgical interventions, during the revision, a complete anatomical rupture of the spinal cord was found in 9 patients, in 27 cases, a picture of hematomyelia and spinal cord contusion at the level of damage. Conclusions: 1 in the presence of symptoms of spinal cord injury, it is necessary to perform le, revision of the spinal canal and spinal cord. 2, the most reliable fixation is achieved when using a construct for tpf, which makes it possible to activate the patients early.

Keywords: trauma, spine, spinal cord, revision, metal structure, tpf system.

Relevance: according to domestic and foreign authors, the treatment of patients with spinal injuries remains an urgent problem of modern neurosurgery to date. The importance of this task is determined by the ever-increasing number of patients with severe spinal injury. Making a decision on surgical intervention on an injured spine without neurological deficit is a difficult task and causes ambiguous judgments.

The transition from conservative to early surgical methods for the treatment of severe spinal injuries led to a significant reduction in disability and a decrease in negative consequences.

Objective of the study: to improve the results of surgical treatment of injuries and diseases of the spine using tpf systems.

Materials and methods: from 2010 to 2020, 114 patients with injuries of varying severity of the thoracic and lumbar spine were under our supervision in the department of neurosurgery of the af rncemmp. The age of patients is from 16 to 60 years. The duration of spinal injury was from 3 hours to 5 days. When distributing patients by gender: men - 76 patients and women - 38. According to the mechanism of injury: falling from a height - 81 patients, road injury - 33 patients.

Examination of patients was carried out according to the approved protocol (examination by specialists, radiography of the spine in standard projections, computed and magnetic resonance imaging of the spine).



Fig.1. Fracture of the body of Tn12 vertebra III-IV degree with posterior displacement

Among the examined patients with complicated spinal injuries, there were 40 patients and 74 patients with uncomplicated injuries. According to the level of the damaged spine: Th 7 - Th 8 - 6 patients, Th 9 - Th 10 - 19 patients, Th 11 - Th 12 - 49 patients, Th12 - L1 - 26 patients, L1 - L2 - 11 patients and L3 vertebra - 3 sick. As can be seen from the above data, the largest number of patients was with injuries of the transitional spine - 86 patients. According to the number of damaged vertebrae: at the level of 1 vertebra - 87 patients and at the level of 2 vertebrae - 27 patients. According to the degree of compression of the vertebral bodies, I degree - 36 patients, II degree - 49 patients, III degree - 16 patients, and with IV degree of damage were - 3 patients. Of the total number of examined patients, 53 patients underwent conservative treatment and 61 patients underwent surgical treatment.

After examinations, 61 patients underwent the following types of surgical interventions:

1. Fixation of the damaged segment of the spine with various types of metal structures (TPF) - 22 patients.
2. Decompressive laminectomy with revision of the spinal canal and spinal cord at the level of injury (without fixation of the spinal column) – 20 patients.
3. Decompressive laminectomy at the level of damage + revision of the epi- and subdural spaces of the spinal cord and, as the final stage of the operation, fixation with metal structures - 19 patients.




Fig.2. Stabilization of the damaged segment of the spine with the TPF system.

Of the 61 patients, 12 used metal structures from the Medbiotech company (Republic of Belarus) and in 29 cases TPF structures from the ChM company (Republic of Poland).

Results: During the surgical interventions, during the revision, a complete anatomical rupture of the spinal cord was found in 9 patients, in 27 cases, a picture of hematomyelia and spinal cord contusion at the level of damage.

During repeated examinations of patients after 3 and 6 months, a satisfactory functioning of the fixing metal structures was noted, after 1 year, looseness of the structures was found in 6 patients, in these cases operations were performed with the re-establishment of the structure and in 2 cases their removal.

Conclusions: Thus, based on the analysis of the results of surgical treatment of patients with spinal injuries, it can be concluded that the most reliable fixation is achieved when using a construction for TPF, which makes it possible to activate patients early.

BIBLIOGRAPHY:

1. Aganesov A.G., Meskhi K.T., Kheilo A.L. Surgical treatment of spinal injuries // 9th Congress of Traumatologists and Orthopedists of Russia: Proceedings. report Saratov, 2010, p. 567.
2. Borzykh K.O., Roerich V.V., Rakhmatillaev Sh.N. Surgical treatment of uncomplicated burst fractures of the thoracic and lumbar vertebrae, accompanied by a critical displacement of fragments into the spinal canal. report Saratov, 2010, p. 585
3. Vetrile S.T., Kuleshov A.A., Shvets V.V. Surgical treatment of fractures of the thoracic and lumbar spine with the use of modern technologies // 9th congress of traumatologists and orthopedists of Russia: Proceedings. report Saratov, 2010. pp. 596-597.
4. Gaidar B.V., Dulaev A.K., Orlov V.P. Surgical treatment of patients with injuries of the thoracic and lumbar spine // Surgery of the spine. 2004. No. 3. S. 40-45.
5. Kelmakov V.P. Complex neurosurgical treatment of patients with post-traumatic spinal cord cysts associated with spinal canal deformity: Abstract of the thesis. dis. ... cand. honey. Sciences. Novosibirsk, 2005.
6. Ramikh E.A. Injuries of the thoracic and lumbar spine // Surgery of the spine. 2008. No. 1. S. 86-106.
7. Roerich V.V., Borzykh K.O., Rakhmatillaev Sh.N. Surgical treatment of burst fractures of the thoracic and lumbar vertebrae, accompanied by narrowing of the spinal canal. 2007. No. 2. S. 8-15.
8. Cigliano A, Scarano E, De Falco R, et al. The posterolateral approach in the 9. Dai LY. Remodeling of the spinal canal after thoracolumbar burst fractures. ClinOrthopRelat Res. 2001;(382):119-123.
9. Denis F. The three column spine and its significance in the classification of acute thoracolumbar spinal injuries. Spine. 1983;8(8):817-831.
10. Morrison RH, Thierolf A, Weckbach A. Volumetric changes of iliac crest autografts used to reconstruct the anterior column in thoracolumbar fractures: a follow-up using CT scans. Spine. 2007;32(26):3030-3035.
11. Mumford J, Weinstein JN, Spratt KF, et al. Thoracolumbar burst fractures. The clinical efficacy and outcome of nonoperative management. Spine. 1993;18(8):955-970.
12. Razak M, Mahmud M, Mokhtar SA, et al. Thoracolumbar fracture-dislocation results of surgical treatment. Med J Malaysia. 2000;55(Suppl. C):14-17.
13. Zdeblick TA, Sasso RC, Vaccaro AR, et al. Surgical treatment of thoracolumbar fractures. Instr. Course Lect. 2009;58:639-644.