

---

# Laparoscopic Methods in the Diagnosis and Treatment of Emergency Surgical Diseases of the Abdominal Cavity

---

*Nurboboyev Adhambek Uygunovich, Makhmudova Guljamol Fazliddinovna, Sokhibova Ziyoda Rakhmonovna*

*Assistants Bukhara State Medical Institute*

---

**Abstract:** The creation of qualitatively new laparoscopic equipment, the improvement of laparoscopic instruments required the revision of outdated provisions and the development of modern principles of laparoscopic surgery. In 2018 – 2020, 20493 operations were performed in the Bukhara branch, of which 9461 (12.1%) were cavity operations. The examined group consisted of 2,479 (26.2%) patients with urgent diseases of the abdominal cavity, among all the abdominal operations performed in the period from 2018 to 2020 using laparoscopic technologies.

**Keywords:** laparoscopic equipment, abdominal surgery, abdominal trauma, fibroesophagoduodenoscopy.

---

**Relevance:** Despite the wide arsenal of non-invasive research methods, the diagnosis of acute surgical diseases of the abdominal cavity in some cases is very difficult. In this situation, the use of laparoscopy allows timely diagnosis and determination of surgical tactics. A comprehensive analysis of the use of laparoscopic operations in emergency abdominal surgery and their improvement is an urgent task of modern clinical surgery. With the development of laparoscopic surgery and the accumulation of experience, the interest of many surgeons in the use of this method in emergency surgery is expanding. Within the framework of general scientific and technological progress, the constant introduction of new technologies makes them an integral part of a particular field of medicine. Today, emergency surgery is impossible to imagine without laparoscopy. In many complex cases, laparoscopy remains indispensable, as it allows you to give a direct visual assessment of the pathological process, identify its prevalence, perform a targeted biopsy and perform instrumental palpation. It is safe to say that the number of urgent operations performed by laparoscopic method is steadily increasing. This circumstance requires the development of clear recommendations on the use of laparoscopic technologies in emergency surgery. The method makes it possible to carry out diagnostics with a greater degree of informativeness, and when establishing pathology in the abdominal cavity, it provides an opportunity to transform diagnostic manipulation into therapeutic. In the Clinic of Surgical Diseases and Resuscitation of the Bukhara State Medical Institute, from 2018 to 2020, out of 9461 emergency abdominal operations performed, laparoscopic technique was used in 2479 (26.2%) observations.

The following laparoscopic operations were performed: laparoscopic cholecystectomy (LCE) for acute calculous cholecystitis was performed in 1984 (80%) patients, diagnostic and therapeutic laparoscopy -49 (1.9%), diagnostic and therapeutic thoracoscopy in 32 (1.3%) patients, laparoscopic appendectomy -18 (0.7%), suturing of perforative ulcers of the stomach and duodenum (duodenum) -15(0.6%)patients, laparoscopic cystectomy (gynecology)-

112(4.5%), laparoscopic tubectomy -181(7.3%), laparoscopic dissection of adhesions - 24 (0.9%) and laparoscopic ureterolithotomy - 64(2.6%) patients.

**The purpose of the study:** Improving the results of surgical treatment of patients with acute surgical diseases of the abdominal cavity, by improving diagnostic and therapeutic laparoscopy. **Materials and methods of research:** The work was carried out in the Bukhara branch of the Emergency Hospital on the basis of surgical departments. In 2018 – 2020, 20493 operations were performed in the Bukhara branch, of which 9461 (12.1%) were cavity operations. The examined group consisted of 2,479 (26.2%) patients with urgent diseases of the abdominal cavity, among all the abdominal operations performed in the period from 2018 to 2020 using laparoscopic technologies.

The paper analyzes the results of a parallel comparative study of laparoscopic and open methods of surgical treatment. All the admitted patients were subjected to a comprehensive clinical study according to the generally accepted scheme using traditional laboratory and radiological methods, general clinical and biochemical studies were conducted; in-depth study of the coagulation system; overview radiography of the abdominal cavity, radiopaque studies of the gastrointestinal tract (gastrointestinal tract); computed tomography (CT), including with contrast, ultrasound examination (ultrasound) of the abdominal cavity and pelvis, magnetic resonance imaging in rare cases - to clarify the diagnosis; endoscopic studies fibroesophagogastroduodenoscopy. According to the indications, patients with cholelithiasis underwent endoscopic retrograde pancreato-cholangiography, supplemented, if necessary, by endoscopic papillosphincterotomy. The patients subjected to laparoscopic examination and surgical treatment were aged from 23 to 77 years, while there were 961 patients older than 60 years (38.7%). The following laparoscopic operations were performed: laparoscopic cholecystectomy (LCE) for acute calculous cholecystitis was performed in 1984 (80%) patients, diagnostic and therapeutic laparoscopy - 49 (1.9%), diagnostic and therapeutic thoracoscopy in 32 (1.3%) patients, laparoscopic appendectomy - 18 (0.7%), suturing of perforative ulcers of the stomach and duodenum (DPC) - 15 (0.6%) patients, laparoscopic cystectomy (gynecology) - 112 (4.5%), laparoscopic tubectomy -181 (7.3%), laparoscopic dissection of adhesions - 24 (0.9%) and laparoscopic ureterolithotomy - 64 (2.6%) patients.

Table No. 1

Types of laparoscopic operations in the Bukhara branch of the Emergency Hospital 2018-2020yy

Operation by years	2018	2019	2020	2018-2020
Diagnostical and treatment of laparoscopy	12	21	16	49(1,9%)
Appendectomy via the Laparoscopy	5	6	7	18(0,7%)
Cholisisectomia via the Laparoscopy	603	708	673	1984(80%)
Laparoscopy in the Perforative ulcer	4	5	6	15(0,6%)
Laparoscopic dissection of adhesions	4	8	12	24(0,9%)
Laparoscopic cystectomy (gynecology)	35	30	47	112(4,5%)
Laparoscopic tubectomy	72	74	35	181(7,3%)
Diagnostic and therapeutic thoracoscopy	2	8	22	32(1,3%)
Ureterolithotomy			64	64(2,6%)
Total laparoscopic operations	737	860	882	2479

To assess the effectiveness of the technique of surgical intervention, the following indicators were evaluated: the duration of surgery, normalization of body temperature, the need for drug stimulation of the intestine, the recovery time of intestinal peristalsis, the time of activation of the patient, the volume of intraoperative blood loss, the duration of hospitalization, the need

and duration of stay in the intensive care unit (ICU), the need and duration of the use of analgesics, the number and the severity of the intra- and postoperative complications, cosmetic defect after surgery, the need for repeated surgical interventions, the number of conversions.

Laparoscopic intervention was performed in a limited hermetic space (in the abdominal, thoracic cavity), which was created by the introduction of gas. A certain level of intra-abdominal pressure (12-15 mmHg) was created by introducing carbon dioxide into the abdominal cavity and was stably maintained throughout the operation, which provided favorable conditions for performing the necessary manipulations under the control of a laparoscope. The gas was injected through a Veresh needle by puncture of the anterior abdominal wall. The air cushion obtained in the abdominal cavity prevented damage to its organs during the introduction of trocars. A Veresh needle was inserted into the abdominal cavity through a small incision (up to 1 cm) in the umbilical region along the white line of the abdomen. To prevent damage to internal organs or vessels with a Veresh needle, the anterior abdominal wall was gripped with a brush into a longitudinal fold or a clamp and lifted up. After the creation of the pneumoperitoneum, trocars were inserted into the abdominal cavity, starting with the paraumbilical, through which a laparoscope was inserted into the abdominal cavity. Other trocars were administered under visual laparoscopic control. Laparoscopic instruments were introduced through trocars, with the help of which laparoscopic intervention was performed.

### **Results and discussions:**

As a result of 1984 laparoscopic cholecystectomies (LCE), complications of LCE for acute cholecystitis were observed in 14 (0.8%) patients. Parenchymal bleeding from the gallbladder bed was observed in 11 (0.6%) patients; in all, bleeding was stopped by laparoscopic coagulation. Damage to the extrahepatic bile ducts occurred in 3 (0.2%) patients and required laparotomy. In 2 observations, a lateral lesion of the choledochus was sutured. Completely crossed hepaticocholedoch 1 patients were restored on the drainage of the Kerus.

Damage to the bile ducts is facilitated by various anatomical variants of the structure and mutual disposition of the tubular structures of the hepatoduodenal ligament. An extremely dangerous situation is when the common hepatic duct is attached directly to the neck of the gallbladder, and the latter has a short duct. We have repeatedly encountered similar cases, but in one of them we could not avoid a complete intersection of the common bile duct.

Damage to the hepaticocholedocha can also occur in the absence of anatomical prerequisites, if the technique of performing the operation is violated. These are the so-called "classic" cases of duct injury. They occur when excessive traction of the gallbladder is created. In order to prevent complications in the isolation of the cystic duct and artery, we use the following technique: using an electric hook, we dissect the peritoneum from the lateral side of the gallbladder to the level of the confluence of the cystic duct into the common hepatic duct, then we isolate the medial wall of the gallbladder with mandatory tissue preparation behind the cystic duct. Thus, the shape of the neck is created according to the type of "elephant trunk", which makes it easy to differentiate the cystic duct to the confluence with the common hepatic duct. If damage to the hepaticocholedochus is detected during surgery, it is advisable to apply its primary suture with a duct width of at least 5 mm with a sufficient wall thickness. Reconstructive surgery requires an experienced surgeon, the necessary tools and suture material.

In the postoperative period, complications occurred in 22 (1.1%) patients: bile drainage from the abdominal cavity - in 8 (0.4%), subhepatic abscess in the bed of the gallbladder — in 2 (0.1%), suppuration of epigastric puncture was observed — in 12 (0.6%) patients. (Table 2.)

Table 2. Complications after laparoscopic cholecystectomy

Types of complications	Number of abs patients (%)
Bile drainage from the abdominal cavity	8 (0,4%)
Subhepatic abscess in the bed of the gallbladder	2 (0,1%)
Suppuration of the epigastric puncture	12 (0,6%)
Total complications	22 (1,1%)8

Laparoscopic interventions made it possible to reduce the length of stay of patients in the hospital (by 22.2 - 37.5%), reduce the percentage of unnecessary laparotomies (by 15-25%), shorten the time of diagnosis, eliminate the occurrence of postoperative hernias, reduce the appointment of narcotic analgesics, transfer patients to enteral nutrition at an earlier date.

Conclusion: Within the framework of general scientific and technological progress, the constant introduction of new technologies makes them an integral part of a particular field of medicine. The current trend in the development of surgery is to reduce the invasiveness and traumaticity of surgical interventions. Due to the development of surgical technologies, one of the main tasks in surgery is to minimize surgical trauma and the subsequent reduction in the number of postoperative complications and mortality, as well as the duration of inpatient treatment of patients while maintaining the quality of surgical care. Today, urgent surgery can no longer be imagined without laparoscopy.

#### List of literature

1. Abdullaev M.A. The causes of conversion during laparoscopic operations in patients with perforated gastroduodenal ulcers / M.A. Abdullaev, Yu.V. Plotnikov, A.M. Mukhanna // Almanac of the A.V. Vishnevsky Institute of Surgery. - 2010. - Vol. 5, No. 1. - p. 71.
2. Abu R.N. Application possibilities video laparoscopic interventions in diagnosis and treatment in emergency abdominal surgery / Ukrainian Scientific and Medical Youth Journal. - 2013. - No. 3. - pp. 15-18.
3. Adamyan A.I. Laparoscopic cholecystectomy in acute cholecystitis: dis. ... Candidate of Medical Sciences: 14.00.27 - Surgery / Research Institute of SP named after N.V. Sklifosovsky. - M., 1997. - 267 p.
4. Alexandrov K.R. Laparoscopic operations from a single access using standard instruments / K.R. Alexandrov, R.A. Burya, I.Y. Illarionov [et al.] // Almanac of the A.V. Vishnevsky Institute of Surgery. -2011. - Vol. 6, No. 1. - pp. 34-35.
5. Akhmedova Maftun Amin kizi. Ko'rak bezi-o'ziga hos intracranial a'zo // Journal of Advanced Research and Stability (JARS) // Volume: 01.05/2021., 171-180 bet. Rakhmonovna, S. Z., & Sharipovna, A. N. (2020). Characteristics of exchange of essential microelements of copper and zinc in healthy fertilized women and women with combined copper and zinc deficiency state. *European Journal of Molecular & Clinical Medicine*, 7(1), 3332-3335.
6. Turdiyev, M. R., Teshayev Sh. J. Morphometric Assessment of Functional Immunomorphology of White Rat Spleen in the Age Aspect *American Journal of Medicine and Medical Sciences* 2019, 9(12): 523-526

7. AT Cho'liyev., US Mamedov., MA Akhmadova., RR Navro'zov., DF Narziyeva Diagnostics of exinococcosis in youth at the modern stage./Journal of Natural Remedies.2021,№1(1).-P37-40
8. Guljamol Fazliddinonvna Makhmudova, Adkhambek Uygunovich Nurboboyev. Treatment of mechanical jaundice via the modern way// Scientific progress, 2021.-№6.-P.530-537
9. Makhmudova G.F. Age-related clinical,anatomical and morphological features of malignant tumors of the cervix// Journal of science and technology//2021.-P.-475-480
10. Turdiev M.R., Teshaev Sh.J. Comparative characteristics of the morphological and morphometric parameters of the spleen of white rats in normal conditions, chronic radiation sickness and correction with a biostimulant // Problems of biology and Medicine - 2020. - No. 4 (120) - P. 160-165
11. AH Narzulloyevich, MG Fazliddinovna, KF Sharopovna// Comparison of the results of modern methods of treatment of elderly women with breast cancer// Eurasian Medical Research Periodical 3, 9-15.
12. Махмудова Г. Ф., Темирова, Д. В., &Баротова, Ш. Б. (2021). Бачадон бўйни хавфли ўсмаларининг ёшга хосхусусиятлари//Academicresearchineducationalsciences // 2(5).-Б.-186-196. <https://doi.org/10.24411/2181-1385-202100871>
13. Makhmudova G.F.,Soxibova Z.R., Mamedov U.S., Nurboboyev A.U. Fertil va keksa yoshli ayollarda bachadon bo'yni xavfli o'smalari tahlili (Buxoro viloyatida)//Oriental Renaissance: Innovative, educational, natural and social sciences//2021.-V 8.-B. 175-184.
14. Nurboboyev A.U., Makhmudova G.F. Miniinvazive approach in the complex treatment of tumor and stone etiology of mechanical jaundice// International journal on Orange technology// Vol 3. Issue 9. Sep.2021.-P. 85-90
15. М.А. Ахмадова, А.Т.,Сохибова З.Р., Д.К. Худойбердиев.,Ж.Р.Нуров Диагностика эхинококкоза у молодёжи на современном этапе./Тиббиётда янги кун 2019 й.3(27)-стр 54-56
16. М.А. Ахмадова, А.Т. Чўлиев, Ж.Р. Нуров, Д.К. ХудойбердиевЛучеваядиагностикаэхинококкозапечени./Биологияватиббиётмуаммолари.2019,№4.2(115)с.20-25
17. Сохибова З.Р.,Ахмадова М.А. Комплексная диагностика и хирургическое и хирургическое лечение осложненных форм эхинококкоза печени./Oriental Renaissance: Innovative, Educational, natural and social sciences/2021й -стр 203-212.
18. НарзиеваД.Ф.ЗначениеИммуногистохимических маркеров при метастазировании рака молочной железы в легкие.// Oriental Renaissance: Innovtive, educational, natural and social sciences.// -2021 Vol.1-C.170-175
19. Abdullayev Habibulla Narzulloyevich, Makhmudova Guljamol Fazliddinovna, Makhmudova Anora Fazliddinovna // Age-related clinical and instrumental analysis of malignant tumors of the cervix// Eurasian Medical Research Periodical.-2021 Vol 3, 1-8.
20. Z.R. Sokhibova, M.R. Turdiyev, (2021). Some Features Of Laboratory Indicators Of Micro And Macro-Elementary Condition Of The Organism Of Female Age Women Innormality And In Iron Deficiency. *The American Journal of Medical Sciences and Pharmaceutical Research*, 3(02), MO- 145.

21. Mamedov U.S., Pulatova D.SH. The Results of Cancer Treatment of the Oral Caviti Tumors in //the Republic of Uzbekistan European journal of Pharmaceutical and Medical Research. -2019. - 6(9). - P. 326-329.
22. Narziyeva D.F., Jonibekov J.J.; Morphological features of tumor in different treatment options for patients with locally advanced breast cancer // Middle European scientific bulletin. Volume 7- 2020-Dec. – P. 105-10
23. Nurov Jamshid Raxmatovich. Morph functional characters of the greater omentum // International Journal of Discoveries and Innovations in Applied Sciences. – 2021. – Vol. 1(5). – P. 130-134.
24. Nurov J.R., Khalikova F.S. Long-term results of surgical treatment patients with stomach cancer // Вестник науки и образования. – 2020. – №23-2(101). – С. 85-89.
25. Р.Р.Наврұзов. Характеристика морфометрических параметров желудка белой крысы в раннем постнатальном периоде // Новый день в медицине. 2 (34/3) 2021 С.17-23
26. Р.Р.Наврұзов. Морфологические и морфометрические изменения слоя желудка месячных белых крыс // Журнал Фогинновационных разработок в фармацевтической и технической науке (JIDPTS). Объем:4, Выпуск:5, Май:2021 стр. :(7-10)
27. R. R. Navruzov. Morphofunctional features of the lymphoid structures of the colon in normal and under the influence of a biostimulator on the background of radiation sickness // Web of Scientist: International Scientific Research Journal Sep 8, 2021 Page: (53-56)
28. Р. Р. Наврұзов., Тешаев Ш.Ж., Очилов К.Р., Худойбердиев Д.К. Сравнительная характеристика толстой кишки белых беспородных крыс при хронической лучевой болезни и после воздействия биостимулятора асд-2ф // Новый день в медицине 6 (38) 2021г. С. 272-276
29. Гафур Нормуродович Саидов, Учкун Гафурович Абдукаримов, Гулжамол Фазлиддиновна Махмудова. Эпидемиологические показатели первично-множественных опухолей (обзор литературы)// Биология и интегративная медицина// 2019№ 11 (39).-С.
30. Нуров Ж.Р. Послеоперационная аналитика раннего периода хирургического лечения злокачественной опухоли желудка // Oriental Renaissance: Innovative, educational, natural and social sciences. – 2021. – Vol. 1(8). – P. 185-191.
31. Sokhibova Z.R., Akhmedova M.A. Complex diagnostics and surgical and surgical treatment of complicated forms of liver echinococcosis./Oriental Renaissance: Innovative, Educational, natural and social sciences/2021 -pp. 203-212.
32. Nurov Jamshid Raxmatovich, Narziyeva Dilnoza Fakhriddinovna. The Significance of Immunohistochemical Markers in the Treatment of Breast Cancer // International journal on orange technology. – 2021. – Vol. 03(9). – P. 69-72.
33. Nurov Jamshid Raxmatovich, Ahmadova Maftuna Amin qizi. Features of Anatomy of the Greater Omentum // International journal on orange technology. – 2021. – Vol. 03(9). – P. 66-68.
34. Nurov Jamshid Raxmatovich, Narziyeva Dilnoza Fakhriddinovna. Immediate Results of Surgical Treatment of Gastric Cancer // International journal on orange technology. – 2021. – Vol. 03(9). – P. 62-65.

35. Sokhibova, Z. R., & Turdiyev, M. R. (2021). Some Features Of Laboratory Indicators Of Micro And Macro-Elementary Condition Of The Organism Of Female Age Women Innormality And In Iron Deficiency. *The American Journal of Medical Sciences and Pharmaceutical Research*, 3(02), 140-145.
36. MG Fazliddinova, NA Uygunovich, ND Faxriddinova The modern way of diagnosis of cervical pathology in women with uterine fibroids via the colposcopy//Web of scientist: international scientific research journal.-3(02), 1017-1027, 2022.
37. Mamedov U.S, Khalikova F. Sh. Advantages of Magnetic Resonance Computer Tomography in the Diagnosis of Thyroid Cancer //Pindus Journal of Culture, Literature, and ELT. – 2021. – Т. 9. – С. 80-84.
38. Axmedov Farhod Hakimovich// Central Asian journal of medical and natural science// Морфологические Изменения Внутри И Внепеченочных Протоков, И Сфинктеров У Больных С Желчекаменной Болезнью, Постхолецистэктомии. Volume: 02 Issue: 05 | Sep-Oct 2021
39. Xudoyberdiyev Dilshod Karimovich characteristics o morphometric parameters of the white rat's stomach in the early postnatal period// Тиббиётда янги кун// 2 (34/3) 2021 С-17-23
40. Xudoyberdiyev Dilshod Karimovich МОРФОЛОГИЧЕСКИЕ И МОРФОМЕТРИЧЕСКИЕ ИЗМЕНЕНИЯ СТЕНКИ ЖЕЛУДКА ОДНОМЕСЯЧНЫХ БЕЛЫХ КРЫС// INTERDISCIPLINARY RESEARCH: SCIENTIFIC HORIZONS AND PERSPECTIVES International Scientific and Theoretical Conference// March 12, 2021 С 57-61
41. Axmedov Farhod Hakimovich SCIENTIFIC COLLECTION «INTERCONF» COMPARATIVE MORPHOMETRY OF INTRA AND EXTRAHEPATIC BILIARY TRACT, BILIARY SPHINCTERS IN PATIENTS WITH CHOLELITHIASIS WHO UNDERWENT CLASSICAL AND LAPAROSCOPIC CHOLECYSTECTOMY № 78 | October, 2021 P-325-327
42. Sultanova L. The Dj.Nuraliyev N.A. Indicators of Seeding of Microorganisms translated from the Large Intestine to Internal Oragan under the Influence of Acuteationation in the Experiment//American Journal of Medicine and Medical Sciences//Volume 10, Number 11,Novembr 2020.p-929-932
43. G.F.Makhmudova Colposcopic analysis of cervical pathology in women with uterine fibroids//Scientific progress// 3(1), 289-296,2022
44. А.У. Нурбобоев, МС Шаропова, А.Ф. Махмудова Турли этиологияли механик сарикликни даволашда замонавий минилапаратом усуллар// Scientific progress// 3(1), 713-721, 2022