
Morphometric Estimates of Histological Structures of the Spleen of White Rats in Postnatal Ontogenesis

M. R. Turdiev, Z. R. Sohibova
Bukhara State Medical Institute

Abstract: This article is devoted to an urgent problem – the development of the spleen as an organ of the immune system at the stages of postnatal ontogenesis. In an experiment on 40 white mongrel male rats of 5 age groups using histological and morph metric methods followed by statistical analysis, data on the morph metric parameters of the spleen of white rats in postnatal ontogenesis are presented. It was found that the studied quantitative parameters of the white pulp of white rats undergo significant postnatal changes, expressed in different age aspects.

Keywords: rats, spleen, white pulp, age-related changes, morph metric parameters.

Relevance. The immune system of humans and animals is one of the most sensitive systems of the body, which quickly reacts to external influences [6].

The largest peripheral organ of the immune system is the spleen [2; 5; 9], one of the main functions of which is the formation of a generalized immune response to the effects of various pathogenic agents, which ensures the maintenance of immune homeostasis and, consequently, the necessary level of adaptive potential of the body [3; 5,14,15].

The immunocompetent compartment of the spleen parenchyma is represented by its white pulp (WP), where two main B- and T-dependent zones are formed, that is, areas of localization of mainly B- and T-lymphocytes. These are, respectively, lymphoid follicles (LF) and per arterial lymphoid couplings (PALM) [2; 4; 5; 8].

A large number of scientific studies based on the study of this organ as a person have been devoted to the study of reactive changes of the spleen, in particular the structural elements of WP [4;5;8;12,17], and various agricultural and laboratory animals [1;3;4;7,16].

Despite the existence of numerous modern studies on the structure of the spleen under the influence of external factors [6], the issues of morph functional changes in the lymphoid formations of the spleen in the age aspect remain insufficiently studied. To accurately represent the reactions of the lymphoid structures of the spleen to any external stimuli or foreign elements, it is necessary to know the morphology of the limophid structures of the spleen normally.

The purpose of the study. To study the regularities of the age dynamics of the features of the histological structure of the white pulp of the spleen of white mongrel rats in postnatal ontogenesis from birth to 12 months of age.

Materials and methods.

The study was conducted on 40 white male mongrel rats, weighing from 20 to 280 g., which were kept in vivarium conditions with a standard diet, free access to water, and the usual mode of purification. The slaughter of rats was carried out under ether anesthesia at the ages

of newborns, 90, 180, 270 and 360 days from the beginning of the experiment. The rats were divided into five age groups, 8 individuals each: group 1 - newborns, group 2 - 90-day-old rats, group 3 - 180-day-old rats, group 4 - 270-day-old rats, group 5 - 360-day-old rats. The studies were carried out in compliance with the rules of humane treatment of animals, which are regulated by the "Rules for carrying out work using experimental animals" approved by the Ethics Committee of the Bukhara State Medical Institute named after Abu Ali ibn Sino (No. 18 of 16.01.2018), and were also based on the provisions of the Helsinki Declaration of the World Medical Association of 1964, amended in 1975, 1983, 1989, 1996, 2000, 2002, 2004, 2008, 2013 years .

For morphological examination, splenic fragments were fixed in a 10% formalin solution, passed through a battery of alcohols and poured into paraffin blocks according to generally accepted methods. Paraffin sections 5-8 microns thick were stained with hematoxylin - eosin. Histological and morphometric examination was performed using a Leica CME microscope, with a Leica Microsystems CH-9435 digital camera. At the same time, the relative area of the white pulp (in%), the diameters of lymphoid nodules (in microns), germinative centers (in microns), the width of the mantle zone (in microns), the marginal zone (in microns) of lymph nodes were determined on the sections of the tear.

Mathematical processing was performed directly from the general data matrix "Excel 7.0" with the involvement of the capabilities of the program "STTGRAPH 5.1", the standard deviation and representativeness errors were determined.

Results and their discussion.

In newborn white rats, the spleen is covered from the outside with a thin connective tissue capsule, from which trabeculae extend deep into the organ. The white pulp of the spleen in newborn white rats is just beginning to form. It is quite difficult to distinguish between per arterial lymphatic couplings and lymphoid nodules, mantle and marginal zones can be determined in the formed nodules.

The study of histological preparations of other age groups of the spleen of white rats allowed us to establish certain relationships between its structural and functional zones. The spleen of white rats has a well-defined connective tissue capsule and trabeculae containing blood vessels.

The parenchyma of the spleen of white rats is represented by red and white pulp. The composition of the red pulp includes sinusoidal capillaries and splenic cords. The white pulp consists of numerous lymph nodes and per arterial lymphoid couplings.

The lymphoid tissue of the spleen, which forms a white pulp, on the sections represents rounded or elongated gray areas among the red pulp filled with erythrocytes. The white pulp is a cluster of lymphocytes around an eccentrically located arterial canal.

The relative area of the white pulp of the spleen of newborn white rats was $17.16 \pm 1.06\%$, the greatest increase in this indicator was observed at 3 months of age ($22.2 \pm 1.72\%$), and the largest at 12 months of age ($12.32 \pm 1.08\%$). After 3 months of age, there is a decrease in this indicator. The diameter of lymph nodes increases by 1.92 times before the age of 3 months. After 3 months of age, this indicator also gradually decreases, where at 12 months of age it is equal to 407.98 ± 8.18 microns. Germinative centers of lymph nodes in newborn white rats with histological sections have not been detected, the greatest increase in this indicator is observed at 3 months of age (147.8 ± 7.66 microns), and the largest at 12 months (120.02 ± 5.36 microns). The width of the mantle zone increases by 1.32 times before the age of 6 months. After 6 months of age, this indicator also gradually decreases, where at 12 months of age it is equal to 41.32 ± 1.12 microns. The width of the marginal zone and the per

arterial zone from newborn age in accordance with age aspects began to increase to 6 months of age, starting from 6 months to one year of age decreased slightly.

The white pulp of the spleen of 3-month-old white rats is represented by well-defined lymph nodes and per arterial lymphoid couplings, among the lymph nodes are dominated by nodules with germinative centers. In 3-month-old white rats, the width of the per arterial zone prevails over other indicators. In 9 and 12-month-old white rats, invasive changes prevail in lymph nodes and per arterial lymphatic couplings.

The percentage of white pulp in newborn white rats is quite low. According to our data, the proportion of white pulp increased markedly in subsequent age groups. The largest values of the white pulp area were observed in the spleen of white rats at 3 months of age, in other age groups, the size of the white pulp steadily decreased, confirming the fact of age-related organ involution.

In terms of the area of the white pulp, the diameter of the lymph nodes and the diameter of the germinal centers, the greatest growth is observed at 3 months of age, and in terms of the width of the mantle zone, the width of the marginal zone and the width of the per arterial zone, the greatest growth is observed at 180 days of age.

With age, all the studied morphometric parameters gradually decrease and the lowest indicator is noted at 12 months of age.

Conclusions. In the process of postnatal ontogenesis, the spleen of white rats undergoes structural and functional transformations reflecting the processes of formation of white pulp. In newborn white rats, the spleen is functionally immature, with hard-to-distinguish zones and poorly formed lymphoid nodules. At three and nine months of age, the spleen has the maximum potential for immunogenesis. At one year of age, the processes of age-related organ involution begin, which manifest themselves in a change in the germinal centers of lymphoid nodules, a decrease in the number and area of lymphoid nodules and later the width of the mantle, marginal, per arterial zone in the white pulp of the spleen.

Literatures

1. Andreeva S. D. characteristics of lymphoid structures of the pig spleen // Mezhdunar. Journal. applied and foundation. research. 2013, No. 4, pp. 97-98.
2. Afanasyev Yu. I., Yurina N. A., Kotovsky E. F. [et al.]. Histology, Embryology, cytology : textbook / 6th ed., rework. and add. / 2012. - 800 p.
3. Bashina S. I. new in the method of studying the spleen of a large pig White breeds // Vestn. Bryansk State University. agricultural enterprise. akad — - 2013. - № 2. - pp. 28-29.
4. Volkov V. P. new algorithm of morphometric assessment of functional immunomorphology of the spleen // Universum: Medicine and pharmacology: electron. scientific. Journal. 2015. № 5-6 (18) .
5. Zaitsev V. B., Fedorovskaya N. S., Diakonov D. A. [et al.] Morph functional characteristic of the human spleen // morphology. 2013, No. 3, pp. 27-31.
6. Kashchenko S. A., Bobrysheva I. V. features of histological structure white pulp of the rat spleen in different periods of postnatal ontogenesis in the conditions of experimental immunosuppression. Medical University. 2014, No. 1, pp. 51-54.
7. Turdiev M.R., Teshayev Sh.J. Comparative characteristics of the morphological and morphometric parameters of the spleen of white rats in normal conditions, chronic

- radiation sickness and correction with a biostimulant // Problems of biology and Medicine - 2020. - No. 4 (120) - P. 160-165
8. Statsenko E. A. modern ideas about the anatomy of the spleen a person // in Ukrainian. med. ALM. — 2009. - Vol. 12, № 3. - pp. 229-232.
 9. Shepitko V. I., Katsai V. V., Stetsuk E. V. Morphofunctional characteristics of the normal spleen of rats // world of Medicine and biology. —2010. - №3. - pp. 99-100
 10. Di Ianni M., Papa B.D., Terenzi A. Interleukin 7- engineered stromal celss a new approach for hastening naïve T-cell recruitment // Hum. Gen. Ther. – 2005. -V.16, №6 - P.752-754) Bertrand J.Y. Fetal spleen stroma drives macrophage commitment / Julien Y.
 11. Bertrand, Guillaume E. Desanti, Richard Lo-Man, Claude Leclerc, Ana Cumano and Rachel Golub // Development. — 2006. — № 133. — P. 3619-3628.
 12. Pereira S.A., Rodrigues D.B., Castro E.C. [et al.] Morphometric study of the spleen in chronic Chagas' disease // Am. J. Trop. Med. Hyg. — 2002. — V. 66, N. 4. — P. 401—403.
 13. Wolber F. M. Roles of spleen and liver in development of the murine hematopoietic system. / Wolber, F. M., Leonard, E., Michael, S., Orschell-Traycoff, C. M., Yoder, M. C. and Srour, E. F. // Exp. Hematol. — 2002. — №30. — 1010-1019.
 14. Turdiev M.R., Teshayev S.J. Comparative characteristics of the spleen of white rats in normal and chronic radiation sickness // Chief Editor. T. 7. P. 11.
 15. Turdiyev, M. R., Teshayev Sh. J. Morphometric Assessment of Functional Immunomorphology of White Rat Spleen in the Age Aspect American Journal of Medicine and Medical Sciences 2019, 9(12): 523-526
 16. Turdiyev, M. R., & Sokhibova, Z. R. (2021). Morphometric Characteristics Of The Spleen Of White Rats In Normal And In Chronic Radiation Disease. The American Journal of Medical Sciences and Pharmaceutical Research, 3(02), 146-154.
 17. Sokhibova, Z. R., & Turdiyev, M. R. (2021). Some Features Of Laboratory Indicators Of Micro And Macro-Elementary Condition Of The Organism Of Female Age Women Innormality And In Iron Deficiency. The American Journal of Medical Sciences and Pharmaceutical Research, 3(02), 140-145.
 18. Nurboboyev A.U., Makhmudova G.F. Miniinvasive approach in the complex treatment of tumor and stone etiology of mechanical jaundice// International journal on Orange technology// Vol 3. Issue 9. Sep.2021.-P. 85-90.
 19. Makhmudova G.F. Age-related clinical, anatomical and morphological features of malignant tumors of the cervix// Journal of science and technology//2021.-P.-475-480.
 20. Rakhmonovna, S. Z., & Sharipovna, A. N. (2020). Characteristics of exchange of essential microelements of copper and zinc in healthy fertilized women and women with combined copper and zinc deficiency state. *European Journal of Molecular & Clinical Medicine*, 7(1), 3332-3335.
 21. Sokhibova Z.R., Akhmedova M.A. Complex diagnostics and surgical and surgical treatment of complicated forms of liver echinococcosis./Oriental Renaissance: Innovative, Educational, natural and social sciences/2021 -pp. 203-212.
 22. Sultanova L. The Dj. Nuraliyev N.A. Indicators of Seeding of Microorganisms translated from the Large Intestine to Internal Oragans under the Influence of Acuteationation in the Experiment//American Journal of Medicine and Medical Sciences//Volume 10, Number

11,Novembr 2020.p-929-932

23. М.А. Ахмадова, А.Т.,Сохибова З.Р., Д.К. Худойбердиев.,Ж.Р.Нуров Диагностика эхинококкоза у молодёжи на современном этапе./Тиббиётда янги кун 2019 й.3(27)-стр 54-56
24. Abdullayev Habibulla Narzulloyevich, Makhmudova Guljamol Fazliddinovna, Makhmudova Anora Fazliddinovna// Age-related clinical and instrumental analysis of malignant tumors of the cervix// Eurasian Medical Research Periodical.-2021 Vol 3, 1-8.
25. Mamedov U.S., Pulatova D.SH. The Results of Cancer Treatment of the Oral Caviti Tumors in //the Republic of Uzbekistan European journal of Pharmaceutical and Medical Research. -2019. - 6(9). - P. 326-329.
26. Narziyeva D.F., Jonibekov J.J.; Morphological features of tumor in different treatment options for patients with locally advanced breast cancer // Middle European scientific bulletin. Volume 7- 2020-Dec. – P. 105-10
27. Nurov Jamshid Raxmatovich. Morph functional characters of the greater omentum // International Journal of Discoveries and Innovations in Applied Sciences. – 2021. – Vol. 1(5). – P. 130-134.
28. R. R. Navruzov. Morph functional features of the lymphoid structures of the colon in normal and under the influence of a biostimulator on the background of radiation sickness // Web of Scientist: International Scientific Research Journal Sep 8, 2021 Page: (53-56)
29. MG Fazliddinovna, NA Uygunovich, ND Faxriddinovna The modern way of diagnosis of cervical pathology in women with uterine fibroids via the colposcopy//Web of scientist: international scientific research journal.-3(02), 1017-1027, 2022.
30. Axmedov Farxod Hakimovich// CENTRAL ASIAN JOURNAL OF MEDICAL AND NATURAL SCIENCES// Морфологические Изменения Внутри И Внепеченочных Протоков, И Сфинктеров У Больных С Желчекаменной Болезнью, Постхолецистэктомии. Volume: 02 Issue: 05 | Sep-Oct 2021
31. Xudoyberdiyev Dilshod Karimovich CHARACTERISTICS OF MORPHOMETRIC PARAMETERS OF THE WHITE RAT'S STOMACH IN THE EARLY POSTNATAL PERIOD// Тиббиётда янги кун// 2 (34/3) 2021 С-17-23
32. G.F.Makhmudova Colposcopic analysis of cervical pathology in women with uterine fibroids//Scientific progress// 3(1), 289-296,2022